

SOLENOZETES GALLONAE SP. NOV., FIRST RECORD OF THE PLASMOBATIDAE IN AUSTRALIA (ACARI: ORIBATIDA)

GLENN S. HUNT

Hunt, G.S. 1994 06 01: *Solenozetes gallonae* sp. nov., first record of the family Plasmobatidae in Australia (Acari: Oribatida). *Memoirs of the Queensland Museum* 35(1): 129-134. Brisbane. ISSN 0079-8835.

Solenozetes gallonae sp. nov. is described from rainforest in Queensland, Australia, and a rediagnosis given for *Solenozetes* Grandjean, 1932 (Plasmobatidae). The status and distribution of other species in the genus are summarised and generic level characters analysed.

□ Acari, Oribatida, Plasmobatidae, *Solenozetes*, Australia, rainforest.

Glenn S. Hunt, Division of Invertebrate Zoology, Australian Museum, PO Box A285, Sydney South, New South Wales 2000, Australia, 15 June 1993.

The Plasmobatidae Grandjean, 1961 is a small group of oribatid mites with a mostly tropical-subtropical distribution extending from the Oriental Region and western Pacific to the eastern Palaeartic and Neotropical Regions. They inhabit litter, moss and epiphytes on tree trunks.

Together with the Hermanniellidae, they constitute the superfamily Hermannielloidea, a taxon characterised by distinctive lateral tubes which bear the pore of the lateral opisthosomal glands (gla in Fig. 1D). Adult Plasmobatidae carry the exuviae ("scalps") of the larva and of three nymphal instars (Fig. 1A,B) which obscure the inconspicuous notogastral setae. They are readily distinguished from the Hermanniellidae which carry a tightly appressed and inconspicuous tritonymphal scalp which bears conspicuous setae. The gnathosoma of the Plasmobatidae is characterised by an elongate subcapitulum above which lie the elongate chelicerae. Definitions of both families are given by Grandjean (1962).

The Plasmobatidae comprises three genera, *Plasmobates* Grandjean, 1929, *Solenozetes* Grandjean, 1932 and *Orbiculobates* Grandjean, 1961, which may be distinguished by the key in Balogh & Balogh (1988, 1992). Generic level characters are analysed below with insights provided by *Solenozetes gallonae* sp. nov. The occurrence of this species in Australia is an important addition to the known distribution of the family.

MATERIALS AND METHODS

In general, terminology follows Balogh & Balogh (1992), and Grandjean (1961) and Norton (1977) for leg chaetotaxy. The shapes of setae and surface ornamentation follow Mahunka & Zombori (1985). Measurements are in micrometres,

those for body length were made ventrally from the tip of the rostrum to the rear of the scalps. A Cambridge Stereoscan 120 was used for SEM. Repositories for type material are AM, Australian Museum, Sydney; ANIC, Australian National Insect Collection, Canberra; QM, Queensland Museum, South Brisbane.

CHARACTER ANALYSIS AT GENERIC LEVEL

Scalps. Grandjean (1932) erected *Solenozetes* as he considered the type species, *Plasmobates cribratus* Grandjean, 1929 from Venezuela, possessed only a tritonymphal scalp. In his 1961 paper, however, he regarded this as an error (scalps can fall off) and questioned the separate status of *Solenozetes*. He nevertheless retained the genus, without giving a clear rediagnosis. Apart from an anterior indentation which characterises the *Orbiculobates* tritonymphal scalp (Grandjean, 1961), scalp morphology appears to be useful only at the species level.

Shape of lateral tubes. Balogh (1972) used this as a key character to distinguish *Plasmobates* from *Solenozetes* and was followed, rather tentatively, by Mahunka (1983) when describing *S. flagellifer* from Mexico. This key character was abandoned by Balogh & Balogh (1988; 1992) and appears to be of value only at the species level.

Number of genital setae and presence/absence of aggenital setae. Balogh & Balogh (1988) transferred *P. carinatus* Hammer, 1961 from Peru and *P. flagellatus* Balogh & Mahunka, 1969 from Amazonia to *Solenozetes* on the basis that they had 6 pairs of genital setae and no aggenital setae (*Plasmobates* has 7 pairs and 1 pair respectively).




FIG. 1. *Solenozetes gallonae* sp. nov. A,C,D,F, frontal. A, whole body; C, prodorsum; D, lateral tube with cerotegument; F, rostral area. B, whole body, lateral; E, notogaster without scalps, dorsal; G, detail lateral tube area without cerotegument and scalps (bo, bothridium; gla, lateral tube of notogastral gland; ia, fissura ia; in, interlamellar seta; ro, rostral seta; ss, sensillus). Scale bars: A,B,E, 100 μ m; C, 50 μ m; D,F,G, 20 μ m.

P. hyalinus Hammer, 1971 from Fiji, *P. acutirostrum* Hammer, 1973 from Western Samoa and *P. asiaticus* Aoki, 1973 from Japan have 6 pairs of genital and no aggenital setae. Hammer (1979) describes *P. javensis* from Java as having 7 pairs of long genital setae but no aggenital setae. I have examined the type and it appears to have 6 genital setae on the right side and a doubtful alveolus of a seventh on the left. However, formal placement of some or all of these species in *Solenozetes* should await a thorough revision of the family including reappraisal of characters and a re-examination of the type of *S. cribratus*.

P. schubarti Perez-Inigo & Baggio, 1988 from Brazil has 7 pairs of genital setae and 1 pair of aggenital setae and would presumably remain in *Plasmobates* together with the type species, *P. pagoda* Grandjean, 1929 from Martinique.

Shape of bothridium. Except perhaps for *O. transvectus* (Grandjean, 1929), descriptions of species with 6 pairs of genital setae (and *P. javensis*), seem to indicate that the bothridial wall forms a partial hood over the bothridial cavity but is deeply invaginate anterolaterally (Fig. 1C). In *P. pagoda* and *P. asiaticus* (7 pairs), the bothridium is not hooded and has a much more open structure. It is invaginate in the mesad as well as the laterad wall. A "hooded" bothridium may be a shared derived character for *Solenozetes* and *Orbiculobates*, as suggested by its absence in *Hermaniellidae* and *Lioididae*, outgroups which have a low, open bothridium. The more open bothridium seems to be correlated with a shorter sensillus.

Notogastral sculpturing. Most *Plasmobates* and *Solenozetes* have a pitted (alveolate) notogaster. The two *Orbiculobates* species, *O. orbiculus* (Grandjean, 1929) and *O. transvectus*, both from Martinique, have a smooth notogaster with scalp retention assisted by an anterior tuft of filaments. *Orbiculobates australis* Balogh & Csizsar, 1963 from Argentina, however, has a smooth notogaster without filaments. It has subsequently been placed in *Solenozetes* by Balogh & Balogh (1988).

Accessory tooth on movable finger of chelicera (as in Fig. 2C). Grandjean (1929: pl. 3F; 1961) does not record it but R.A. Norton (pers. comm.) reports it in a new species of *Orbiculobates* and several probably undescribed *Solenozetes* spp. and *Plasmobates* spp. It is present in *S. gallonae* and on the types of *S. carinatus* and *P. javensis*. The structure appears to be a shared

derived character at the family level. Its function is unknown but it may assist in breaking food.

SYSTEMATICS

Solenozetes Grandjean, 1932

Solenozetes Grandjean, 1932: 653; 1961: 127; Balogh, 1972: 57; Balogh & Balogh, 1988: 87; 1992: 42.

TYPE SPECIES

Plasmobates cribratus Grandjean, 1929; by monotypy.

DIAGNOSIS

Plasmobatidae with 6 pairs of genital setae but no aggenital setae. Anterior margin of notogaster without mesial tuft of filaments and correlated invagination of tritonymphal scalp. Notogaster usually alveolate or foveolate; lateral tubes of variable shape. Sensillus very long, filiform or sometimes flagellate; posterior wall of bothridium forms partial hood over bothridial cavity, anteriolateral wall deeply excavate allowing sensillus greater movement in laterad direction. Chelicera of at least some species with strong accessory tooth on movable finger.

Solenozetes gallonae sp. nov.

(Figs 1-3)

ETYMOLOGY

For Ms Julie Gallon who collected the material.

TYPE MATERIAL

HOLOTYPE: adult, QM, Queensland, foot of Blackbutt Range, 5 km. E. of Benarkin, rainforest, leaf litter, QM berlesate 522, J. Gallon, 30 Mar 1983.

PARATYPES: QM, same data, 2 adults; AM, same data, 1 adult; ANIC, same data, 1 adult.

OTHER MATERIAL

QMS20116 (SEM Stub No. 142), foot of Blackbutt Range, 5 km. E. of Benarkin, rainforest, leaf litter, QM berlesate 522, J. Gallon, 30 Mar 1983, 2 adults; QMS20117 (SEM Stub No. 100), same data, 5 adults; QM (SEM Stub No. 129), same data, adult notogaster without scalps.

DIAGNOSIS

Sensillus much longer than distance between interlamellar setae, filiform, not flagellate, and not expanding subdistally; lateral tubes directed only slightly forward and distally expanded into




FIG. 2. *Solenozetes gallonae* sp. nov. A-G, ventral aspect. A, gnathosoma; B, epimeral region; C, right chelicera; D, whole animal; E, genital plates; F, pedipalp; G, anal plates. a,m,h, gnathosomal setae; ad3, adanal seta; at, accessory tooth; b, lateral boss between legs I and II; sp, sword-like process. Scale bars: A,B,E, 50µm; D, 100µm; C, 10µm; F,G, 20µm.

ly, but eupathidic nature of setae p and s not resolved. Ontogeny of setae not studied. Size order of subcapitular setae seems to reflect anteriad hypertrophy of the subcapitulum.

ACKNOWLEDGEMENTS

This research is supported by an Australian Biological Resources Survey grant. I thank Dan Bickel, Australian Museum, Bruce Halliday, C.S.I.R.O. and Roy Norton, State University of New York, for commenting on the manuscript, and Henrik Enghoff of the University Zoological Museum, Copenhagen for the loan of Hammer's *Plasmobates* types. Suzanne Lindsay took the SEMs, Ross Pogson assisted with polarising microscopy and Roger Springthorpe did the camera ready layouts.

LITERATURE CITED

- AOKI, J. 1973. Oribatid mites from Iriomote-jima, the southernmost island of Japan. *Memoirs of the National Museum of Tokyo* 6:85-101.
- BALOGH, J. 1972. 'The oribatid genera of the world'. (Akademiai Kiado: Budapest).
- BALOGH, J. & BALOGH, P. 1988. Oribatid mites of the neotropical region I. In Balogh, P. (ed.), 'The soil mites of the world 2'. (Elsevier: Amsterdam).
1992. 'The oribatid mites genera of the world', Vol. I. (Hungarian Natural History Museum: Budapest).
- BALOGH, J. & CSISZAR, J. 1963. The zoological results of Gy. Topal's collectings in Argentina. 5. Oribatei (Acarina). *Annales Historico-Naturales Musei Nationalis Hungarici* 55: 463-485.
- BALOGH, J. & MAHUNKA, S. 1969. The zoological results of the Hungarian Soil Zoological Expeditions to South America 11. Acari: oribatids from the material of the Second Expedition. II. *Opuscula Zoologica* 9(1): 31-68.
- GRANDJEAN, F. 1929. Quelques nouveaux genres d'Oribatei du Venezuela et de la Martinique. *Bulletin de la Societe Zoologique de France* 54: 400-423.
1932. Observations sur les oribates (2e serie). *Bulletin Museum National d'Histoire Naturelle, Paris, 2nd series*, 1931, 3(7): 651-665.
1961. Les Plasmobatidae n. fam. (Oribates). *Acarologia* 3(1): 96-129.
1962. Au sujet des Hermanniellidae (Oribates). *Premiere Partie. Acarologia* 4(2): 237-273.
- HAMMER, M. 1961. Investigations on the oribatid fauna of the Andes Mountains II. Peru. *Biologiske Skrifter udgivet af det Kongelige Danske Videnskabernes Selskab* 13(1): 1-150.
1971. On some oribatids from Viti Levu, the Fiji Islands. *Biologiske Skrifter udgivet af det Kongelige Danske Videnskabernes Selskab* 16(6): 1-56.
1973. Oribatids from Tongatapu and Eua, the Tonga Islands, and from Upolu, Western Samoa. *Biologiske Skrifter udgivet af det Kongelige Danske Videnskabernes Selskab* 20(3): 4-70.
1979. Investigations on the oribatid fauna of Java. *Biologiske Skrifter udgivet af det Kongelige Danske Videnskabernes Selskab* 22(9): 1-78.
- MAHUNKA, S. 1983. Neue und interessante Milben aus dem Genfer Museum XLV. *Oribatida Americana* 6: Mexico II (Acari). *Revue Suisse Zoologie* 90(2): 269-298.
- MAHUNKA, S. & ZOMBORI, L. 1985. The variability of some morphological features in oribatid mites. *Folia Entomologica Hungarica* 46(1): 115-128.
- NORTON, R.A. 1977. A review of Grandjean's system of leg chaetotaxy in the Oribatei and its application to the Damaeidae. Pp. 33-62. In Dindal, D.L. (ed.), 'Biology of Oribatid Mites'. (State University New York College of Environmental Science and Forestry: Syracuse, New York).
- PEREZ-INIGO, C. & BAGGIO, D. 1988. Oribates edaphiques du Bresil (IV). Oribates de l'Etat de Sao Paulo (Premiere Partie). *Acarologia* 29(2): 189-204.